

Acta Arith. 127(2007), no. 2, 103–113.

MIXED SUMS OF SQUARES AND TRIANGULAR NUMBERS

ZHI-WEI SUN (NANJING)

ABSTRACT. By means of q -series, we prove that any natural number is a sum of an even square and two triangular numbers, and that each positive integer is a sum of a triangular number plus $x^2 + y^2$ for some $x, y \in \mathbb{Z}$ with $x \not\equiv y \pmod{2}$ or $x = y > 0$. The paper also contains some other results and open conjectures on mixed sums of squares and triangular numbers.

1. INTRODUCTION

A classical result of Fermat asserts that any prime $p \equiv 1 \pmod{4}$ is a sum of two squares of integers. Fermat also conjectured that each $n \in \mathbb{N}$ can be written as a sum of three triangular numbers, where $\mathbb{N}$ is the set $\{0, 1, 2, \dots\}$ of natural numbers, and triangular numbers are those integers $t_x = x(x+1)/2$ with $x \in \mathbb{Z}$. An equivalent version of this conjecture states that $8n + 3$ is a sum of three squares (of odd integers). This follows from the following profound theorem (see, e.g., [G, pp. 38–49] or [N, pp. 17–23]).

Gauss-Legendre Theorem. *$n \in \mathbb{N}$ can be written as a sum of three squares of integers if and only if n is not of the form $4^k(8l + 7)$ with $k, l \in \mathbb{N}$.*

Building on some work of Euler, in 1772 Lagrange showed that every natural number is a sum of four squares of integers.

For problems and results on representations of natural numbers by various quadratic forms with coefficients in $\mathbb{N}$, the reader may consult [Du] and [G].

Motivated by Ramanujan's work [Ra], L. Panaitopol [P] proved the following interesting result in 2005.

Theorem A. *Let a, b, c be positive integers with $a \leq b \leq c$. Then every odd natural number can be written in the form $ax^2 + by^2 + cz^2$ with $x, y, z \in \mathbb{Z}$, if and only if the vector (a, b, c) is $(1, 1, 2)$ or $(1, 2, 3)$ or $(1, 2, 4)$.*

2000 *Mathematics Subject Classification*: Primary 11E25; Secondary 05A30, 11D85, 11P99.

Supported by the National Science Fund for Distinguished Young Scholars (Grant No. 10425103) in China.

According to L. E. Dickson [D2, p. 260], Euler already noted that any odd integer $n > 0$ is representable by $x^2 + y^2 + 2z^2$ with $x, y, z \in \mathbb{Z}$.

In 1862 J. Liouville (cf. [D2, p. 23]) proved the following result.

Theorem B. *Let a, b, c be positive integers with $a \leq b \leq c$. Then every $n \in \mathbb{N}$ can be written as $at_x + bt_y + ct_z$ with $x, y, z \in \mathbb{Z}$, if and only if (a, b, c) is among the following vectors:*

$$(1, 1, 1), (1, 1, 2), (1, 1, 4), (1, 1, 5), (1, 2, 2), (1, 2, 3), (1, 2, 4).$$

Now we turn to representations of natural numbers by mixed sums of squares (of integers) and triangular numbers.

Let $n \in \mathbb{N}$. By the Gauss-Legendre theorem, $8n + 1$ is a sum of three squares. It follows that $8n + 1 = (2x)^2 + (2y)^2 + (2z + 1)^2$ for some $x, y, z \in \mathbb{Z}$ with $x \equiv y \pmod{2}$; this yields the representation

$$n = \frac{x^2 + y^2}{2} + t_z = \left(\frac{x + y}{2}\right)^2 + \left(\frac{x - y}{2}\right)^2 + t_z$$

as observed by Euler. According to Dickson [D2, p. 24], E. Lionnet stated, and V. A. Lebesgue [L] and M. S. Réalis [Re] proved that n can also be written in the form $x^2 + t_y + t_z$ with $x, y, z \in \mathbb{Z}$. Quite recently, this was reproved by H. M. Farkas [F] via the theory of theta functions.

Using the theory of ternary quadratic forms, in 1939 B. W. Jones and G. Pall [JP, Theorem 6] proved that for any $n \in \mathbb{N}$ we have $8n + 1 = ax^2 + by^2 + cz^2$ for some $x, y, z \in \mathbb{Z}$ if the vector (a, b, c) belongs to the set

$$\{(1, 1, 16), (1, 4, 16), (1, 16, 16), (1, 2, 32), (1, 8, 32), (1, 8, 64)\}.$$

As $(2z + 1)^2 = 8t_z + 1$, the result of Jones and Pall implies that each $n \in \mathbb{N}$ can be written in any of the following three forms with $x, y, z \in \mathbb{Z}$:

$$2x^2 + 2y^2 + t_z = (x + y)^2 + (x - y)^2 + t_z, \quad x^2 + 4y^2 + t_z, \quad x^2 + 8y^2 + t_z.$$

In this paper we establish the following result by means of q -series.

Theorem 1. (i) *Any $n \in \mathbb{N}$ is a sum of an even square and two triangular numbers. Moreover, if $n/2$ is not a triangular number then*

$$\begin{aligned} & |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \nmid x\}| \\ &= |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \mid x\}|. \end{aligned} \quad (1)$$

(ii) *If $n \in \mathbb{N}$ is not a triangular number, then*

$$\begin{aligned} & |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \not\equiv y \pmod{2}\}| \\ &= |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \equiv y \pmod{2}\}| > 0. \end{aligned} \quad (2)$$

(iii) A positive integer n is a sum of an odd square, an even square and a triangular number, unless it is a triangular number t_m ($m > 0$) for which all prime divisors of $2m + 1$ are congruent to 1 mod 4 and hence $t_m = x^2 + y^2 + t_z$ for some integers $x > 0$ and z with $x \equiv m/2 \pmod{2}$.

Remark. Note that $t_2 = 1^2 + 1^2 + t_1$ but we cannot write $t_2 = 3$ as a sum of an odd square, an even square and a triangular number.

Here are two more theorems of this paper.

Theorem 2. Let a, b, c be positive integers with $a \leq b$. Suppose that every $n \in \mathbb{N}$ can be written as $ax^2 + by^2 + ct_z$ with $x, y, z \in \mathbb{Z}$. Then (a, b, c) is among the following vectors:

$$(1, 1, 1), (1, 1, 2), (1, 2, 1), (1, 2, 2), (1, 2, 4), \\ (1, 3, 1), (1, 4, 1), (1, 4, 2), (1, 8, 1), (2, 2, 1).$$

Theorem 3. Let a, b, c be positive integers with $b \geq c$. Suppose that every $n \in \mathbb{N}$ can be written as $ax^2 + bt_y + ct_z$ with $x, y, z \in \mathbb{Z}$. Then (a, b, c) is among the following vectors:

$$(1, 1, 1), (1, 2, 1), (1, 2, 2), (1, 3, 1), (1, 4, 1), (1, 4, 2), (1, 5, 2), \\ (1, 6, 1), (1, 8, 1), (2, 1, 1), (2, 2, 1), (2, 4, 1), (3, 2, 1), (4, 1, 1), (4, 2, 1).$$

Theorem 1 and Theorems 2–3 will be proved in Sections 2 and 3 respectively. In Section 4, we will pose three conjectures and discuss the converses of Theorems 2 and 3.

2. PROOF OF THEOREM 1

Given two integer-valued quadratic polynomials $f(x, y, z)$ and $g(x, y, z)$, by $f(x, y, z) \sim g(x, y, z)$ we mean

$$\{f(x, y, z) : x, y, z \in \mathbb{Z}\} = \{g(x, y, z) : x, y, z \in \mathbb{Z}\}.$$

Clearly $\sim$ is an equivalence relation on the set of all integer-valued ternary quadratic polynomials.

The following lemma is a refinement of Euler's observation $t_y + t_z \sim y^2 + 2t_z$ (cf. [D2, p. 11]).

Lemma 1. For any $n \in \mathbb{N}$ we have

$$|\{(y, z) \in \mathbb{N}^2 : t_y + t_z = n\}| = |\{(y, z) \in \mathbb{Z} \times \mathbb{N} : y^2 + 2t_z = n\}|. \quad (3)$$

Proof. Note that $t_{-y-1} = t_y$. Thus

$$\begin{aligned}
& |\{(y, z) \in \mathbb{N}^2 : t_y + t_z = n\}| = \frac{1}{4} |\{(y, z) \in \mathbb{Z}^2 : t_y + t_z = n\}| \\
&= \frac{1}{4} |\{(y, z) \in \mathbb{Z}^2 : 4n + 1 = (y + z + 1)^2 + (y - z)^2\}| \\
&= \frac{1}{4} |\{(x_1, x_2) \in \mathbb{Z}^2 : 4n + 1 = x_1^2 + x_2^2 \text{ and } x_1 \not\equiv x_2 \pmod{2}\}| \\
&= \frac{2}{4} |\{(y, z) \in \mathbb{Z}^2 : 4n + 1 = (2y)^2 + (2z + 1)^2\}| \\
&= \frac{1}{2} |\{(y, z) \in \mathbb{Z}^2 : n = y^2 + 2t_z\}| = |\{(y, z) \in \mathbb{Z} \times \mathbb{N} : n = y^2 + 2t_z\}|.
\end{aligned}$$

This concludes the proof. $\square$

Lemma 1 is actually equivalent to the following observation of Ramanujan (cf. Entry 25(iv) of [B, p. 40]): $\psi(q)^2 = \varphi(q)\psi(q^2)$ for $|q| < 1$, where

$$\varphi(q) = \sum_{n=-\infty}^{\infty} q^{n^2} \quad \text{and} \quad \psi(q) = \sum_{n=0}^{\infty} q^{t_n}. \quad (4)$$

Let $n \in \mathbb{N}$ and define

$$r(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n\}|, \quad (5)$$

$$r_0(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \mid x\}|, \quad (6)$$

$$r_1(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n \text{ and } 2 \nmid x\}|. \quad (7)$$

Clearly $r_0(n) + r_1(n) = r(n)$. In the following lemma we investigate the difference $r_0(n) - r_1(n)$.

Lemma 2. *For $m = 0, 1, 2, \dots$ we have*

$$r_0(2t_m) - r_1(2t_m) = (-1)^m(2m + 1). \quad (8)$$

Also, $r_0(n) = r_1(n)$ if $n \in \mathbb{N}$ is not a triangular number times 2.

Proof. Let $|q| < 1$. Recall the following three known identities implied by Jacobi's triple product identity (cf. [AAR, pp. 496–501]):

$$\begin{aligned}
\varphi(-q) &= \prod_{n=1}^{\infty} (1 - q^{2n-1})^2 (1 - q^{2n}) \quad (\text{Gauss}), \\
\psi(q) &= \prod_{n=1}^{\infty} \frac{1 - q^{2n}}{1 - q^{2n-1}} \quad (\text{Gauss}), \\
\prod_{n=1}^{\infty} (1 - q^n)^3 &= \sum_{n=0}^{\infty} (-1)^n (2n + 1) q^{t_n} \quad (\text{Jacobi}).
\end{aligned}$$

Observe that

$$\begin{aligned}
& \sum_{n=0}^{\infty} (r_0(n) - r_1(n)) q^n \\
&= \left(\sum_{x=-\infty}^{\infty} (-1)^x q^{x^2} \right) \left(\sum_{y=0}^{\infty} q^{t_y} \right) \left(\sum_{z=0}^{\infty} q^{t_z} \right) = \varphi(-q) \psi(q)^2 \\
&= \left(\prod_{n=1}^{\infty} (1 - q^{2n-1})^2 (1 - q^{2n}) \right) \left(\prod_{n=1}^{\infty} \frac{1 - q^{2n}}{1 - q^{2n-1}} \right)^2 \\
&= \prod_{n=1}^{\infty} (1 - q^{2n})^3 = \sum_{m=0}^{\infty} (-1)^m (2m+1) (q^2)^{t_m}.
\end{aligned}$$

Comparing the coefficients of q^n on both sides, we obtain the desired result. $\square$

The following result was discovered by Hurwitz in 1907 (cf. [D2, p. 271]); an extension was established in [HS] via the theory of modular forms of half integer weight.

Lemma 3. *Let $n > 0$ be an odd integer, and let $p_1, \dots, p_r$ be all the distinct prime divisors of n congruent to 3 mod 4. Write $n = n_0 \prod_{0 < i \leq r} p_i^{\alpha_i}$, where $n_0, \alpha_1, \dots, \alpha_r$ are positive integers and n_0 has no prime divisors congruent to 3 mod 4. Then*

$$|\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = n^2\}| = 6n_0 \prod_{0 < i \leq r} \left(p_i^{\alpha_i} + 2 \frac{p_i^{\alpha_i} - 1}{p_i - 1} \right). \quad (9)$$

Proof. We deduce (9) in a new way and use some standard notations in number theory.

By (4.8) and (4.10) of [G],

$$\begin{aligned}
& |\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = n^2\}| \\
&= \sum_{d|n} \frac{24}{\pi} d \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{-4d^2}{m} \right) \\
&= \frac{24}{\pi} \sum_{d|n} d \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{2k-1} \sum_{c|\gcd(2k-1, d)} \mu(c) \\
&= \frac{24}{\pi} \sum_{d|n} d \sum_{c|d} \mu(c) \sum_{k=1}^{\infty} \frac{(-1)^{((2k-1)c-1)/2}}{(2k-1)c} \\
&= \frac{24}{\pi} \sum_{d|n} d \sum_{c|d} \frac{\mu(c)}{c} (-1)^{(c-1)/2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{2k-1} \\
&= 6 \sum_{c|n} (-1)^{(c-1)/2} \frac{\mu(c)}{c} \sum_{q|\frac{n}{c}} cq = 6 \sum_{c|n} (-1)^{(c-1)/2} \mu(c) \sigma \left(\frac{n}{c} \right)
\end{aligned}$$

and hence

$$\begin{aligned}
& \frac{1}{6} |\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = n^2\}| \\
&= \sum_{d_0 | n_0} \sum_{d_1 | p_1^{\alpha_1}, \dots, d_r | p_r^{\alpha_r}} \left(\frac{-1}{d_0 d_1 \cdots d_r} \right) \mu(d_0 d_1 \cdots d_r) \sigma \left(\frac{n_0}{d_0} \prod_{0 < i \leq r} \frac{p_i^{\alpha_i}}{d_i} \right) \\
&= \sum_{d_0 | n_0} \left(\frac{-1}{d_0} \right) \mu(d_0) \sigma \left(\frac{n_0}{d_0} \right) \times \prod_{0 < i \leq r} \sum_{d_i | p_i^{\alpha_i}} \left(\frac{-1}{d_i} \right) \mu(d_i) \sigma \left(\frac{p_i^{\alpha_i}}{d_i} \right) \\
&= \sum_{d_0 | n_0} \mu(d_0) \sigma \left(\frac{n_0}{d_0} \right) \times \prod_{0 < i \leq r} \left(\sigma(p_i^{\alpha_i}) + \left(\frac{-1}{p_i} \right) \mu(p_i) \sigma(p_i^{\alpha_i-1}) \right) \\
&= n_0 \prod_{0 < i \leq r} (p_i^{\alpha_i} + 2\sigma(p_i^{\alpha_i-1})) = n_0 \prod_{0 < i \leq r} \left(p_i^{\alpha_i} + 2 \frac{p_i^{\alpha_i} - 1}{p_i - 1} \right).
\end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 1. (i) By the Gauss-Legendre theorem, $4n + 1$ is a sum of three squares and hence $4n + 1 = (2x)^2 + (2y)^2 + (2z + 1)^2$ (i.e., $n = x^2 + y^2 + 2t_z$) for some $x, y, z \in \mathbb{Z}$. Combining this with Lemma 1 we obtain a simple proof of the known result

$$r(n) = |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = n\}| > 0.$$

Recall that $r_0(n) + r_1(n) = r(n)$. If $n/2$ is not a triangular number, then $r_0(n) = r_1(n) = r(n)/2 > 0$ by Lemma 2. If $n = 2t_m$ for some $m \in \mathbb{N}$, then we also have $r_0(n) > 0$ since $n = 0^2 + t_m + t_m$.

(ii) Note that

$$n = x^2 + y^2 + t_z \iff 2n = 2(x^2 + y^2) + 2t_z = (x + y)^2 + (x - y)^2 + 2t_z.$$

From this and Lemma 1, we get

$$\begin{aligned}
& |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n\}| \\
&= |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + 2t_z = 2n\}| \\
&= |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = 2n\}| = r(2n) > 0;
\end{aligned}$$

in the language of generating functions, it says that

$$\varphi(q)\psi(q)^2 + \varphi(-q)\psi(-q)^2 = 2\varphi(q^2)^2\psi(q^2).$$

Similarly,

$$\begin{aligned}
& |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \equiv y \pmod{2}\}| \\
&= |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + 2t_z = 2n \text{ and } 2 \mid x\}| \\
&= |\{(x, y, z) \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N} : x^2 + t_y + t_z = 2n \text{ and } 2 \mid x\}| = r_0(2n) > 0
\end{aligned}$$

and

$$|\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = n \text{ and } x \not\equiv y \pmod{2}\}| = r_1(2n). \quad (10)$$

If n is not a triangular number, then $r_0(2n) = r_1(2n) = r(2n)/2 > 0$ by Lemma 2, and hence (2) follows from the above.

(iii) By Theorem 1(ii), if n is not a triangular number then $n = x^2 + y^2 + t_z$ for some $x, y, z \in \mathbb{Z}$ with $2 \mid x$ and $2 \nmid y$.

Now assume that $n = t_m$ ($m > 0$) is not a sum of an odd square, an even square and a triangular number. Then $r_1(2t_m) = 0$ by (10). In view of (ii) and (8),

$$\begin{aligned} & |\{(x, y, z) \in \mathbb{Z} \times \mathbb{Z} \times \mathbb{N} : x^2 + y^2 + t_z = t_m\}| \\ &= r_0(2t_m) + r_1(2t_m) = r_0(2t_m) - r_1(2t_m) = (-1)^m(2m+1). \end{aligned}$$

Therefore

$$\begin{aligned} (-1)^m(2m+1) &= \frac{1}{2} |\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + t_z = t_m\}| \\ &= \frac{1}{2} |\{(x, y, z) \in \mathbb{Z}^3 : 2(2x)^2 + 2(2y)^2 + (2z+1)^2 = 8t_m + 1\}| \\ &= \frac{1}{2} |\{(x, y, z) \in \mathbb{Z}^3 : (2x+2y)^2 + (2x-2y)^2 + (2z+1)^2 = 8t_m + 1\}| \\ &= \frac{1}{2} |\{(x_1, y_1, z) \in \mathbb{Z}^3 : 4(x_1^2 + y_1^2) + (2z+1)^2 = (2m+1)^2\}| \\ &= \frac{1}{6} |\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = (2m+1)^2\}|. \end{aligned}$$

Since

$$(-1)^m 6(2m+1) = |\{(x, y, z) \in \mathbb{Z}^3 : x^2 + y^2 + z^2 = (2m+1)^2\}| \not\geq 6(2m+1),$$

by Lemma 3 the odd number $2m+1$ cannot have a prime divisor congruent to 3 mod 4. So all the prime divisors of $2m+1$ are congruent to 1 mod 4, and hence

$$|\{(x, y) \in \mathbb{N}^2 : x > 0 \text{ and } x^2 + y^2 = 2m+1\}| = \sum_{d \mid 2m+1} 1 > 1$$

by Proposition 17.6.1 of [IR, p. 279]. Thus $2m+1$ is a sum of two squares of positive integers. Choose positive integers x and y such that $x^2 + y^2 = 2m+1$ with $2 \mid x$ and $2 \nmid y$. Then

$$8t_m + 1 = (2m+1)^2 = (x^2 - y^2)^2 + 4x^2 y^2 = 8t_{(x^2 - y^2 - 1)/2} + 1 + 16 \left(\frac{x}{2}\right)^2 y^2$$

and hence

$$t_m = t_{(x^2-y^2-1)/2} + \left(\frac{x}{2}y\right)^2 + \left(\frac{x}{2}y\right)^2.$$

As $x^2 = 2m + 1 - y^2 \equiv 2m \pmod{8}$, m is even and

$$\frac{m}{2} \equiv \left(\frac{x}{2}\right)^2 \equiv \frac{x}{2} \equiv \frac{x}{2}y \pmod{2}.$$

We are done. $\square$

3. PROOFS OF THEOREMS 2 AND 3

Proof of Theorem 2. We distinguish four cases.

Case 1. $a = c = 1$. Write $8 = x_0^2 + by_0^2 + t_{z_0}$ with $x_0, y_0, z_0 \in \mathbb{Z}$, then $y_0 \neq 0$ and hence $8 \geq b$. Since $x^2 + 5y^2 + t_z \neq 13$, $x^2 + 6y^2 + t_z \neq 47$ and $x^2 + 7y^2 + t_z \neq 20$, we must have $b \in \{1, 2, 3, 4, 8\}$.

Case 2. $a = 1$ and $c = 2$. Write $5 = x_0^2 + by_0^2 + 2t_{z_0}$ with $x_0, y_0, z_0 \in \mathbb{Z}$. Then $y_0 \neq 0$ and hence $5 \geq b$. Observe that $x^2 + 3y^2 + 2t_z \neq 8$ and $x^2 + 5y^2 + 2t_z \neq 19$. Therefore $b \in \{1, 2, 4\}$.

Case 3. $a = 1$ and $c \geq 3$. Since $2 = x^2 + by^2 + ct_z$ for some $x, y, z \in \mathbb{Z}$, we must have $b \leq 2$. If $b = 1$, then there are $x_0, y_0, z_0 \in \mathbb{Z}$ such that $3 = x_0^2 + y_0^2 + ct_{z_0} \geq c$ and hence $c = 3$. But $x^2 + y^2 + 3t_z \neq 6$, therefore $b = 2$. For some $x, y, z \in \mathbb{Z}$ we have $5 = x^2 + 2y^2 + ct_z \geq c$. Since $x^2 + 2y^2 + 3t_z \neq 23$ and $x^2 + 2y^2 + 5t_z \neq 10$, c must be 4.

Case 4. $a > 1$. As $b \geq a \geq 2$ and $ax^2 + by^2 + ct_z = 1$ for some $x, y, z \in \mathbb{Z}$, we must have $c = 1$. If $a > 2$, then $ax^2 + by^2 + t_z \neq 2$. Thus $a = 2$. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $4 = 2x_0^2 + by_0^2 + t_{z_0} \geq b$. Note that $2x^2 + 3y^2 + t_z \neq 7$ and $2x^2 + 4y^2 + t_z \neq 20$. Therefore $b = 2$.

In view of the above, Theorem 2 has been proven. $\square$

Proof of Theorem 3. Let us first consider the case $c > 1$. Since $1 = ax^2 + bt_y + ct_z$ for some $x, y, z \in \mathbb{Z}$, we must have $a = 1$. Clearly $x^2 + bt_y + ct_z \neq 2$ if $c \geq 3$. So $c = 2$. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $5 = x_0^2 + bt_{y_0} + 2t_{z_0} \geq b$. It is easy to check that $x^2 + 3t_y + 2t_z \neq 8$. Therefore $b \in \{2, 4, 5\}$.

Below we assume that $c = 1$. If a and b are both greater than 2, then $ax^2 + bt_y + t_z \neq 2$. So $a \leq 2$ or $b \leq 2$.

Case 1. $a = 1$. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $8 = x_0^2 + bt_{y_0} + t_{z_0} \geq b$. Note that $x^2 + 5t_y + t_z \neq 13$ and $x^2 + 7t_y + t_z \neq 20$. So $b \in \{1, 2, 3, 4, 6, 8\}$.

Case 2. $a = 2$. For some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $4 = 2x_0^2 + bt_{y_0} + t_{z_0} \geq b$. Thus $b \in \{1, 2, 4\}$ since $2x^2 + 3t_y + t_z \neq 7$.

Case 3. $a > 2$. In this case $b \leq 2$. If $b = 1$, then for some $x_0, y_0, z_0 \in \mathbb{Z}$ we have $5 = ax_0^2 + t_{y_0} + t_{z_0} \geq a$, and hence $a = 4$ since $3x^2 + t_y + t_z \neq 8$ and $5x^2 + t_y + t_z \neq 19$. If $b = 2$, then for some $x, y, z \in \mathbb{Z}$ we have $4 = ax^2 + 2t_y + t_z \geq a$ and so $a \in \{3, 4\}$.

The proof of Theorem 3 is now complete. $\square$

4. SOME CONJECTURES AND RELATED DISCUSSION

In this section we raise three related conjectures.

Conjecture 1. *Any positive integer n is a sum of a square, an odd square and a triangular number. In other words, each natural number can be written in the form $x^2 + 8t_y + t_z$ with $x, y, z \in \mathbb{Z}$.*

We have verified Conjecture 1 for $n \leq 15,000$. By Theorem 1(iii), Conjecture 1 is valid when $n \neq t_4, t_8, t_{12}, \dots$.

Conjecture 2. *Each $n \in \mathbb{N}$ can be written in any of the following forms with $x, y, z \in \mathbb{Z}$:*

$$x^2 + 3y^2 + t_z, \quad x^2 + 3t_y + t_z, \quad x^2 + 6t_y + t_z, \quad 3x^2 + 2t_y + t_z, \quad 4x^2 + 2t_y + t_z.$$

Conjecture 3. *Every $n \in \mathbb{N}$ can be written in the form $x^2 + 2y^2 + 3t_z$ (with $x, y, z \in \mathbb{Z}$) except $n = 23$, in the form $x^2 + 5y^2 + 2t_z$ (or the equivalent form $5x^2 + t_y + t_z$) except $n = 19$, in the form $x^2 + 6y^2 + t_z$ except $n = 47$, and in the form $2x^2 + 4y^2 + t_z$ except $n = 20$.*

Both Conjectures 2 and 3 have been verified for $n \leq 10,000$.

The second statement in Conjecture 3 is related to an assertion of Ramanujan confirmed by Dickson [D1] which states that even natural numbers not of the form $4^k(16l + 6)$ (with $k, l \in \mathbb{N}$) can be written as $x^2 + y^2 + 10z^2$ with $x, y, z \in \mathbb{Z}$. Observe that

$$\begin{aligned} n &= x^2 + 5y^2 + 2t_z \text{ for some } x, y, z \in \mathbb{Z} \\ \iff 4n + 1 &= x^2 + 5y^2 + z^2 \text{ for some } x, y, z \in \mathbb{Z} \text{ with } 2 \nmid z \\ \iff 8n + 2 &= 2(x^2 + y^2) + 10z^2 = (x + y)^2 + (x - y)^2 + 10z^2 \\ &\quad \text{for some } x, y, z \in \mathbb{Z} \text{ with } 2 \nmid y \\ \iff 8n + 2 &= x^2 + y^2 + 10z^2 \text{ for some } x, y, z \in \mathbb{Z} \text{ with } x \not\equiv y \pmod{4}. \end{aligned}$$

Below we reduce the converses of Theorems 2 and 3 to Conjectures 1 and 2. For convenience, we call a ternary quadratic polynomial $f(x, y, z)$ *essential* if $\{f(x, y, z) : x, y, z \in \mathbb{Z}\} = \mathbb{N}$. (Actually, in 1748 Goldbach (cf. [D2, p. 11]) already stated that $x^2 + y^2 + 2t_z$, $x^2 + 2y^2 + t_z$, $x^2 + 2y^2 + 2t_z$ and $2x^2 + 2t_y + t_z$ are essential.)

Step I. We show that the 10 quadratic polynomials listed in Theorem 2 are essential except for the form $x^2 + 3y^2 + t_z$ appearing in Conjecture 2.

As $4x^2 + y^2 + 2t_z \sim 4x^2 + t_y + t_z$, the form $x^2 + (2y)^2 + 2t_z$ is essential by Theorem 1(i). Both $x^2 + (2y)^2 + t_z$ and $2x^2 + 2y^2 + t_z = (x + y)^2 + (x - y)^2 + t_z$ are essential by Theorem 1(ii) and the trivial fact $t_z = 0^2 + 0^2 + t_z$. We

have pointed out in Section 1 that $x^2 + 2(2y)^2 + t_z$ is essential by [JP, Theorem 6], and we don't have an easy proof of this deep result.

Since

$$x^2 + 2y^2 + 4t_z \sim x^2 + 2(t_y + t_z) \sim t_x + t_y + 2t_z,$$

the form $x^2 + 2y^2 + 4t_z$ is essential by Theorem B (of Liouville). By the Gauss-Legendre theorem, for each $n \in \mathbb{N}$ we can write $8n + 2 = (4x)^2 + (2y + 1)^2 + (2z + 1)^2$ (i.e., $n = 2x^2 + t_y + t_z$) with $x, y, z \in \mathbb{Z}$. Thus the form $x^2 + 2y^2 + 2t_z$ is essential since $2x^2 + y^2 + 2t_z \sim 2x^2 + t_y + t_z$.

Step II. We analyze the 15 quadratic polynomials listed in Theorem 3.

By Theorem 1(i), $(2x)^2 + t_y + t_z$ and $x^2 + t_y + t_z$ are essential. Since

$$\begin{aligned} x^2 + 2t_y + t_z &\sim t_x + t_y + t_z, \\ x^2 + 2t_y + 2t_z &\sim t_x + t_y + 2t_z, \\ x^2 + 4t_y + 2t_z &\sim t_x + 4t_y + t_z, \\ x^2 + 5t_y + 2t_z &\sim t_x + 5t_y + t_z, \\ 2x^2 + 4t_y + t_z &\sim 2t_x + 2t_y + t_z, \end{aligned}$$

the forms

$$x^2 + 2t_y + t_z, \quad x^2 + 2t_y + 2t_z, \quad x^2 + 4t_y + 2t_z, \quad x^2 + 5t_y + 2t_z, \quad 2x^2 + 4t_y + t_z$$

are all essential by Liouville's theorem. For $n \in \mathbb{N}$ we can write $2n = x^2 + 4t_y + 2t_z$ with $x, y, z \in \mathbb{Z}$, and hence $n = 2x_0^2 + 2t_y + t_z$ with $x_0 = x/2 \in \mathbb{Z}$. So the form $2x^2 + 2t_y + t_z$ is also essential.

Recall that $2x^2 + t_y + t_z$ and $2x^2 + y^2 + 2t_z$ are essential by the last two sentences of Step I. For each $n \in \mathbb{N}$ we can choose $x, y, z \in \mathbb{Z}$ such that $2n + 1 = 2x^2 + (2y + 1)^2 + 2t_z$ and hence $n = x^2 + 4t_y + t_z$. So the form $x^2 + 4t_y + t_z$ is essential.

The remaining forms listed in Theorem 3 are $x^2 + 8t_y + t_z$ and four other forms, which appear in Conjectures 1 and 2 respectively. We are done.

Acknowledgment. The author is indebted to the referee for his/her helpful comments.

Added in proof. The second conjecture in Section 4 has been confirmed by Song Guo, Hao Pan and the author.

REFERENCES

- [AAR] G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Cambridge Univ. Press, Cambridge, 1999.
- [B] B. C. Berndt, *Ramanujan's Notebooks, Part III*, Springer, New York, 1991.
- [D1] L. E. Dickson, *Integers represented by positive ternary quadratic forms*, Bull. Amer. Math. Soc. **33** (1927), 63–70.
- [D2] L. E. Dickson, *History of the Theory of Numbers*, Vol. II, AMS Chelsea Publ., 1999.
- [Du] W. Duke, *Some old problems and new results about quadratic forms*, Notice Amer. Math. Soc. **44** (1997), 190–196.
- [F] H. M. Farkas, *Sums of squares and triangular numbers*, Online J. Anal. Combin. **1** (2006), #1, 11 pp. (electronic).
- [G] E. Grosswald, *Representation of Integers as Sums of Squares*, Springer, New York, 1985.
- [HS] M. D. Hirschhorn and J. Sellers, *On representations of a number as a sum of three squares*, Discrete Math. **199** (1999), 85–101.
- [IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Grad. Texts in Math. 84, Springer, New York, 1990.
- [JP] B. W. Jones and G. Pall, *Regular and semi-regular positive ternary quadratic forms*, Acta Math. **70** (1939), 165–191.
- [L] V. A. Lebesgue, *Questions 1059, 1060, 1061 (Lionnet)*, Nouv. Ann. Math. **11** (1872), 516–519.
- [N] M. B. Nathanson, *Additive Number Theory: The Classical Bases*, Grad. Texts in Math. 164, Springer, New York, 1996.
- [P] L. Panaitopol, *On the representation of natural numbers as sums of squares*, Amer. Math. Monthly **112** (2005), 168–171.
- [Ra] S. Ramanujan, *On the expression of a number in the form $ax^2 + by^2 + cz^2 + du^2$* , in: Collected Papers of Srinivasa Ramanujan, Cambridge Univ. Press, 1927, 169–178.
- [Re] M. S. Réalis, *Scolies pour un théoreme d'arithmétique*, Nouv. Ann. Math. **12** (1873), 212–217.

Department of Mathematics
 Nanjing University
 Nanjing 210093
 People's Republic of China
 zwsun@nju.edu.cn
<http://math.nju.edu.cn/~zwsun>